# Socioeconomic Disparities in Infection-related Hospitalization among the U.S. Home Health Care Population: Insights from a National Study

**Gayani Perera, PhD, MS<sup>1</sup>**; Jingjing Shang, PhD, RN, FAAN, OCN<sup>1</sup>; Jianfang Liu, PhD<sup>1</sup>; Ashley Chastain, DrPH, MPH<sup>1</sup>; Margaret McDonald, MS<sup>2</sup>; David Russell, PhD<sup>2</sup>; Jinjiao Wang, PhD, RN<sup>3</sup> <sup>1</sup>Columbia University School of Nursing, New York, NY; <sup>2</sup>Center for Home Care Policy & Research at VNS Health, New York, NY; <sup>3</sup>University of Rochester School of Nursing, Rochester, NY

## BACKGROUND





- Home health care (HHC) agencies provide skilled nursing care or therapy services to the elderly and acutely ill Medicare beneficiaries within the comfort of their homes.
- About 3.1 million Medicare beneficiaries receive HHC, and infections are a leading cause of hospitalization among HHC patients. Limited evidence has focused on the relationship between the risk of infection and neighborhood socioeconomic status (SES).

Area Deprivation Index (ADI)					
Income	Median Family Income				
	Percent of families below the poverty level				
	Percent of population below 150% of the poverty threshold				
Education	Percent of population age >= 25 years with < 9 years of education				
	Percent of population age >= 25 years with < high school diploma				
Employment	Percent of employed persons aged >= 16 years who are employed in white-collar occupations				
	Percent of civilian labor force population aged >= 16 years who are unemployed				
Housing / Household	Percent of single-parent households with children age < 18 years				
	Percent of households without a motor vehicles				
	Median home value				
	Median gross rent				
	Median monthly mortgage				
	Home ownership rate				
	Percent of households without a telephone				
	Percent occupied housing units without complete plumbing				
	Percent of households more than one person per room ( crowding)				

a more comprehensive measure of ent metrics (Fig. 1) to assess disadvantages at the neighborhood level each neighborhood (at the 9-digit zip into percentiles (based on national scores) from 0 to 100.

Fig. 1 The Area Deprivation Index (ADI) integrates 17 socioeconomic indicators.

### **RESEARCH OBJECTIVE**

To Analyze the association between neighborhood deprivation and infection-related hospitalization of HHC patients prior to the pandemic.

METHODS

Data	<ul> <li>2019 Outcome and Assessment Informati</li> <li>2019 Master Beneficiary Summary File (M</li> <li>2019 Medicare Provider Analysis and Revi</li> <li>2019 Area Health Resources File (AHRF)</li> <li>2019 Area Deprivation Index (ADI)</li> <li>2010 Rural-Urban Commuting Area (RUC</li> <li>60-day HHC episodes were defined from tand linked with hospitalization data (from days of HHC initiation.</li> </ul>
Sample	<ul> <li>3,656,734 Medicare beneficiaries from 8,</li> </ul>
Measurement	<ul> <li>Outcome - Hospitalization due to infection MedPAR ICD-10 principal or admitting dia on admission.</li> <li>Explanatory Variable - Neighborhood de by ADI -categorized into quintiles (Quartil prived, Quintile 5= highest deprived)</li> <li>Covariates - Agency and patient level ch county level health resources.</li> </ul>
Analysis	<ul> <li>Descriptive statistics for overall and by AI</li> <li>Multivariable logistic regression models ac patient, and county-level health resources</li> <li>ADI interaction with sex and dual eligibilit their association with infection-related home.</li> </ul>

• The Area Deprivation Index (ADI) offers neighborhood SES. It includes 17 differbased on 9-digit zip codes. In ADI data, code level) is assigned a score, ranked

> ion Set (OASIS) 4BSF) iew file (MedPAR)

CA) codes. the OASIS MedPAR) within 60

,135 HHC agencies.

on was defined using agnosis codes present

deprivation measured le 1= lowest de-

naracteristics and

DI quintile. djusting for agency,

ty were tested for ospitalizations.

#### Agency (N = 8, 135)Characteristics 18% Quintile 1 Quintile 2 Quintile 3 Quintile 4 20% Quintile 5 18%

**Fig.2** Distribution of (A) HHC Agency (B) HHC Agency Ownership by ADI Quintile.

Compared to patients living in the least deprived neighborhoods (Quintile 1), those living in the most deprived neighborhoods (Quintile 5) were more likely to obtain HHC services from non-profit or government agencies. **Patient Characteristics** 



Fig. 3 Patient-level characteristics across the Area Deprivation Index (ADI).

- Overall, most of HHC patients were White (77.6%) and had an average age of 80. More than half were female (61.7%), and 17.7% resided in rural areas. About 24% of patients were eligible for both Medicare and Medicaid.
- Compared to patients living in the least deprived neighborhoods (Quintile 1), rural areas.

## **County-Level Health Resources**

	Overall	Area Deprivation Index (ADI) Quintile				
		Quintile 1 (1-21%)	Quintile 2 (22-39%)	Quintile 3 (40-57%)	Quintile 4 (58-77%)	Quintile 5 (78-100%)
No. of Hospital Beds						
Zero	3.72	1.91	1.91	3.33	5.18	7.59
>0-512	32.08	9.80	24.58	35.38	44.54	45.58
513-2,473	31.93	31.08	39.48	35.33	28.54	25.61
>2,473	32.26	58.66	34.04	25.96	21.74	21.22
No. of PCPs (%)						
Zero	0.31	0.02	0.04	0.12	0.41	0.94
>0-123	33.32	5.18	19.91	35.75	49.79	55.07
124-740	33.12	27.01	42.86	39.02	30.72	26.45
>740	33.25	67.79	37.19	25.11	19.08	17.54
No. of HHAs (%)						
Zero	8.74	1.51	4.90	8.74	13.00	15.27
>0 to 4	29.83	14.89	24.90	32.00	38.46	38.54
5 to 23	30.87	34.98	35.96	32.67	26.99	24.06
>23	30.56	48.61	34.24	26.58	21.56	22.12

**Table 1** County level characteristics across the Area Deprivation Index (ADI).

Compared to patients living in the least deprived neighborhoods (Quintile 1), those living in the most deprived neighborhoods (Quintile 5) tended to reside in areas with fewer hospital beds, fewer Primary Care Physicians, and HHC agencies (as highlighted in Table 1).



■ Total ■ Quintil 1 ■ Quintile 2 ■ Quintile 3 ■ Quintile 4 ■ Quintile 5

those living in the most deprived neighborhoods (Quintile 5) were less likely to be White or Asian and more likely to be Black, have dual eligibility, and live in



Unadiusted

Adjusted

Sex	ADI Quint	
Male	Quintile 1	
	Quintile 2	
	Quintile 3	
	Quintile 4	
	Quintile S	
Female	Quintile :	
	Quintile 2	
	Quintile 3	
	Quintile 4	
	Quintile !	
Dual Eligibility		
Non-dua	eligibilit	

B

Dual eligibility

Fig. 4 Odds Ratio (OR) of infection-related hospitalization for quintiles of ADI by (A) Unadjusted and Adjusted, (B) ADI and Sex interaction, (C) ADI and Dual Eligibility interaction. CI: Confidence Interval. Quintile 1 is the least deprived (reference), and Quintile 5 is the most deprived. \*\*\* p < 0.01, \*\*p < 0.05, \*p < 0.1



Sex and dual eligibility for Medicare and Medicaid interact with neighborhood deprivation, revealing that females and dual-eligible individuals in socioeconomically disadvantaged areas are particularly at risk for infections.



## COLUMBIA

SCHOOL OF NURSING

## , VNS Health

ROCHESTER MEDICAL CENTER

## Multivariable Logistic Regression Models



## CONCLUSIONS

Residing in the most socioeconomically disadvantaged neighborhoods is associated with an increased risk of infection-related hospitalizations.

## ACKNOWLEDGMENTS

Agency for Healthcare **Research and Quality** 

**R01HS028637** 

**CONTACT INFORMATION** Gayani Perera, PhD Senior Data Manager Center for Health Policy, CUSON uep2000@cumc.columbia.edu